REFERENCES

1. A Effionora, F Nadira formulation and evaluation of phytosome- loaded maltodextrin- gum Arabic microsphere system for delivery of camellia sinesis extract J Young Pharm, 2018; 10(2) 56-62.
2. A Ruggero, C Andrea, S Pasquale, C Giuseppe,R Maria, M Alessandra phytp-liposomes as nanoshuttles for water-insoluble silybin- phospholipids complex, international journal of pharmaceutics volume 471, issue 1-2,2014; 173-181.
3. A semalty, M. Semalty,R.Singh,M S, M. RawatPhytosomes in herbal drug delivery ,Indian drugs, 2006;43:937-946.
4. A Mazumder, A Dwivedi, JLD Preez, JD plessisin vitro wound healing and cytotoxic effects of sinigrin-phytosome complex Int. J Pharm, (2015),498 (1–2): 283-293.
5. Agrawal A.D Pharmacological activities of flavonoids : a review International journal of Pharmaceutical Science and Nanotechnology 2011; 4(2) 1394-1398.
6. Angelico, R. et al. Silybin-phospholipid complex encapsulated into liposomes: preparation, physico-chemical characterization and in vitro test of its antiviral efficiency Int. J. Pharm. 2014, 471, 173.
7. Awasthi R, KulkarniG,Pawar V Phytosomes: an approach to increase the bioavailability of plant extracts https://www.researchgate.net/journal/0975-1491_International_Journal_of_Pharmacy_and_Pharmaceutical_Sciences 2011;3(2) 1-3.
8. Bao V, Sheng Z, Qiao C Flavonoids with therapeutic potential in prostate cancer http://www.eurekaselect.com/node/578 2016; 25: 1205-1229.
9. Beecher G Overview of dietary flavonoids: nomenclature, occurance and intake https://www.ncbi.nlm.nih.gov/pubmed/14519822# 2003;133(10): 3248S-3254S.
10. Bhattacharya S Phytosme: the new technology for enhancement of bioavailability of botanical and nutraceuticals Journal of Aesthetic and Antiaging Medicine 2009; 2(3): 225-232.
11. Bombardelli E, Mustich G (1991) Bilobalide-phospholipid complex, their uses and formulation containing them. U. S. Patent No. EPO-275005.
12. Bonbardelli E, Curri SB, Loggia RD, Negro PD, Tubaro A, Gariboldipcomplexes between phospholipids and vegetal derivatives of biological interest fitoterapia 60(1989), 1-9.
13. Chen A, Xu J, Johnson AC. Curcumin inhibits human colon cancer cell growth by suppressing gene expression of epidermal growth factor receptor through reducing activity of transcription factor Egr-1. https://www.ncbi.nlm.nih.gov/pubmed/16170359# 2006; 25 (2):278–287.

14. Cook N.C &Samman S Flavonoids- chemistry, metabolism, cadioprotectiveeffects,and dietary sources https://www.sciencedirect.com/science/journal/09552863 1996; 7(2): 66-76.
15. Dass MK, Kalita B Phytosme : A Overview Jouranal of pharmaceutical and scientific innovation, 2013 2(3) 7-11.
16. Dhase A, Saboo S Preparation and Evaluation of Phytosomes Containing Methanolic Extract of Leaves of AegleMarmelos (Bael), Int.J. PharmTech Res. 2015,8 (6),pp 231-240.
17. E. Bombardelli, G. Mustich. Bilobalide-phospholipid comlex, their uses and formulation containing them. U. S. Patent No. EPO-275005, 1991.
18. Emerenciano VP, Militao JSLT, Campos CC, Romoff P, Kaplan MAC, Zambon M, Brant AJC: Flavonoids as chemotaxonomic markers for Asteraceae. Biochemical Systematics and Ecology 2001; 29: 947–957.
19. F May, S Wedad, Y Ossama,Abdallahself assembled phospholipids based phytosomalnanocarrieras promising plateforms for improving oral bioavailability of the vanticancercelastrolhttps://www.ncbi.nlm.nih.gov/pubmed/29102699# 2018; 535(1-2) 18-26.
20. Ghanbarzadeh B, Babazadeh A, Hamishehkar H Nano phytosome as a potential food- grade delivery system http://dx.doi.org/10.1016/j.fbio.2016.07.006 2016; 15: 126-135.
21. Ghanbarzadeh B, Babazadeh A, Hamishehkar H. Nano-phytosome as a potential food-grade delivery system. Food Biosci. 2016; 15:126-35.
22. Hao h, jia Y, Han R, IA amp phytosome: an effective approach to enhance the oral bioavailability of active constituents extracted from plants j Chin Pharm sci 22(5) 2013. 385-392.
23. Husch J, Bohnet J, Fricker G, Skarke C, Artaria C, Appendinno G et al.enhancement absorption of bowellic acid by a lecithin delivery from (phytosome) of boswellia extract https://www.ncbi.nlm.nih.gov/pubmed/23092618# 2013;(64): 89-98.
24. J Nilesh, G Brama,TNavneet,jRuchi,BJitendera, j Deepak et al.phytosome : a novel drug delivewry system of herbal medicine internationzal journal of pharmaceutical science and drug research 2010;2(4)-224-228.
25. K Bhupen, D Malay resveratrol- phospholipids complexes (phytosome) with improved physiochemical properties favourable for drug delivery via skin World Journal of Pharmaceutical Research 2015;4(5) 1497-1517.

26. K Govindaranjan, M Mas, K Aminu, A Faridah,FSharida optimization, formulation and characterization of multiflavonoids-loaded flavanosomebybulk or sesquential technique Int. J Nanomedicine 2016; 11:3417-3434.
27. K Junaid, A Alexander, Ajazuddin, S Swarnlata, S Shailendra recent advanced and future prospects of phyto-phopholipidcomplexation technique for improving pharmacokinetics profiles of plant actives, journal of controlled release volume 168, issue 1, 2013;50-60.
28. Kalita B, Das M Rutin–phospholipid complex in polymer matrix for long-term delivery of rutin via skin for the treatment of inflammatory diseasesJournal of artificial cells, nanomedicine, and biotechnology 2018; 46(1) 41-56.
29. Karatas A, Turhan F Phyto-Phospholipid Complexes as Drug Delivery System for Herbal Extracts/Molecules Turk J Pharm Sci 12(1), 93-102, 2015.
30. Kartas A, Turhan F Phyto-phopholipidcomlexescas drug delivery system for herbal extract /molecules turk j pharm sci 12(1), 93-102, 2015.
31. Kidd PM. Bioavailability and activity of phytosome complexes from botanical polyphenols: the silymarin, curcumin, green tea, and grape seed extracts. Altern Med. Rev. 2009;14:226–46.
32. L Ying,Y da, L Shi, C Si-Bao,C Albert process parameters and morphology in puerarin, phospholipids and their complex microparticles generation by supercritical anti-solvent precipitation International Journal of Pharmaceutics 2008; 359: 35–45.
33. Lewandowska U, Szewczyk K, Hrabec E, Janecka A &Gorlach S Overview of metabolism and bioavailability enhancement of polyphenols https://www.ncbi.nlm.nih.gov/pubmed/24295170# 2013; 61(50):12183-12199.
34. M Damle, R MallyaDevelopment and evaluation of a novel delivery system containing phytophospholipid complex for skin aging ,AAPS PharmSciTech, 17 (3) (2016),607-617
35. Mahajan R ,Chaudhari G A novel approach towards phytosomal flavonoids An international journal of pharmaceutical sciences 2012; 3(3): 2079-2105 Kumar S, Pandey A Chemistry and biological activities of flavonoids : an overview The Scientific World Journal 2013; 1-14
36. Mali D, Vyas L, Dwivedi S and Joshi H Investigation on Standardization Parameters of leaves of Thespesiapopulnea Linn. with reference to Formulation and Evaluation of PhytosomesInt. J. of Pharm. Life Sci. 5(6): June, 2014.
37. Mukherjee PK 2001. Evaluation of Indian Traditional Medicine. Drug Information J.; 35(2):623- 631.
38. N Chetan, H KetanPhytosomes- Novel drug delivery system Indian journal of drugs, 2017, 5(1), 16-36.

39. P Yiqiong, Z Xitong, Z Qi, W Bing,CYuxi,ZChuanqi et al. 20 (S)- Protopanaxadiol phospholipid complex: process of optimization , characterization , in Vitro Dissolution and molecular docking studies https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6272886/# 2016;21(10) 1396 or 1-21.
40. Pathan RA, Bhandari U. Preparation & characterization of embelin–phospholipid complex as effective drug delivery tool. J Inclus Phenomena Macrocycl 2011;69(1-2):139-47.
41. Ravi GS, Chandur V, Shabaraya A.R, Sanjay K Phytosomes: an advanced herbal drug delivery system IJPRBS, 2015; 4(3):414-432.
42. Saoji, S.D., Raut, N.A., Dhore, P.W. et al. Preparation and Evaluation of Phospholipid-Based Complex of Standardized *Centella* Extract (SCE) for the Enhanced Delivery of PhytoconstituentsAAPS J (2016) 18(1) 102-114.
43. Shan L, Tan QY, Hong W, Hong L, Zhang jqpreparation, characterization and in vitro anti-tumor activities of evodiamine phospholipids complex Chin Pharm J, 47 (7) (2012), pp. 517-523
44. Sikarwar M, Sharma S, Jain A, Prial S Preparation, characterization and evaluation of marsupsin – phopholipid complex AAPS Pharmscitech.2008; 9(1) 129-137.
45. Singh A, SharanV,Singh M &Bandarin A Phytosome as a drug delivery system for polyphenolicsphytoconstitents Iranian journal of pharmaceutical science 2011; 7(4)209-219.
46. Singh R, Parpani S, Narke R, Chavan R phytosome: recent advance research for novel drug delivery system, Asian Journal of Pharmaceutical Research and Development Vol. 2 (3) May–June. 2014:15-29.
47. Singh RP and Narke R: Preparation and Evaluation of Phytosome of Lawsone. Int. J. Pharm. Sci. Res. 2015; 6(12): 5217-26.
48. Telange, D.R., et al., Formulation and characterization of an apigenin-phospholipid phytosome (APLC) for improved solubility, in vivo bioavailability, an..., European Journal of Pharmaceutical Sciences (2016),1-14.
49. Udapurkar PP, Kamble SR, Bhusnure OG, DiosminPhytosomes: Development, Optimization and Physicochemical Characterization. Indian J of Pharmaceutical Education and Research 2018;52(4S):S29-S36.
50. XuK,Liu B, Ma Y, Du J, Li G, Gao H, et al. Physiochemical properties and antioxidant activities of luteolin- phospholipid complex molecules, 14(9) (2009) 3486-3493.

51. Yu F, Li Y, Chen Q, Yuan H, Wang D. Monodispersemicroparticles loaded with the self-assembled berberinephospholipid complex-based phytosomes for improving oral bioavailability and enhancing hypoglycemic efficiency. Eur J Pharm Biopharm 2016;103:136-48
52. Yue PF, Yuan HL, Li XY, Yang M, Zhu WF, Process optimization, characterization and evaluation in vivo of oxymatrine– phospholipid complex, Int J Pharm 387, 139– 146, 2010.
53. Yue PF, Yuan HL, Li XY, Yang M, Zhu WF. Process optimization, characterization and evaluation *in vivo* of oxymatrine-phospholipid complex. Int J Pharm. 2010;387:139–146
54. Zue W, Yan F, Zhang B, Jiantao L & Mei D Advanced in study of Ginkgo biloba leaves extract on aging related disease, https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5758353/# 2017; 8(6): 812-826.